arXiv:1203.0280v1 [math.GR] 1 Mar 2012

The orbital counting problem for hyperconvex representations

A. Sambarino

Abstract

We show a precise counting result on the symmetric space of a real algebraic non compact semi simple group G, for a representation $\rho : \Gamma \to G$ in the Hitchin component. We also show a mixing property for the Weyl chamber flow in this setting.

Contents

1 Introduction **1**

2 Real Hölder cocycles **6**

3 The action $V \curvearrowright \Gamma\backslash\partial^2\Gamma \times V$ **11**

4 Convex representations and the Weyl chamber flow **17**

5 The orbital counting problem **24**

1 Introduction

The *Orbital Counting Problem* states as following: given a discrete sub group Δ of a connected non-compact real algebraic semi simple Lie group G, find an asymptotic for the growth of

$$\#\{g \in \Delta : d_X(o, g \cdot o) \le t\}$$

when $t \to \infty$ where $o = [K]$ is a point in G's symmetric space $X = G/K$ and K is a maximal compact subgroup of G.

When G has higher rank and the group Δ is a lattice, this problem has been studied by Eskin-McMullen[5]. They show that the number of points on $\Delta \cdot o \cap B(o, t)$ is asymptotic (modulo a constant) to the volume $\mathrm{Haar}(B(o, t))$ of the ball of radius t. Hence, the asymptotic has a polynomial term together with an exponential term. Similar results have been obtained by Duke-Rudnick-Sarnak[4].

The interest is then posed on subgroups of infinite co-volume. An important feature in negative curvature is the limit set of the group on the visual boundary. On higher rank it turns out to be more useful to consider the Furstenberg boundary:

Let P be a minimal parabolic subgroup of G and write $\mathscr{F} = G/P$ for the *Furstenberg boundary* of G's symmetric space. Benoist[1] has shown that a Zariski dense subgroup Δ of G has a smallest closed invariant set for the action $\Delta \curvearrowright \mathscr{F}$, this set is called the *limit set on* $\mathscr{F}$ of Δ and denoted L_Δ .

Schottky groups provide an example of subgroups one can reasonably handle. These are finitely generated free subgroups of G, for which one has good control on the relative position of the fixed points on $\mathscr{F}$ of the generators, together with nice contractions properties.

This precise information allows Quint[13] to build an equivariant continuous function from the boundary of the group (in the sense of Gromov) to Furstenberg's boundary $\mathscr{F}$. The limit set of the Schottky group is thus naturally identified with a sub-shift of finite type. Quint[13] then studies the thermodynamic formalism on this sub-shift and obtains an asymptotic for the orbital counting problem for this class of subgroups. This asymptotic is exponential and does not contain any polynomial term.

Let us introduce the subgroups of G we will study. The product $\mathscr{F} \times \mathscr{F}$ has a unique open G-orbit denoted $\partial^2\mathscr{F}$. For example, when $G = \mathrm{PGL}(d,\mathbb{R})$ the set $\mathscr{F}$ is the set of complete flags of $\mathbb{R}^d$, i.e. families of subspaces $\{V_i\}_{i=0}^d$ such that $V_i \subset V_{i+1}$ and $\dim V_i = i$; and the set $\partial^2\mathscr{F}$ is the set of flags in general position, i.e. pairs $\{V_i\}$ and $\{W_i\}$ such that for every i one has

$$V_i \oplus W_{d-i} = \mathbb{R}^d.$$

Definition 1.1. Let Γ be the fundamental group of a closed negatively curved manifold. We say that a representation $\rho : \Gamma \to G$ is *hyperconvex* if there exists a Hölder continuous equivariant mapping $\zeta : \partial\Gamma \to \mathscr{F}$ such that the pair $(\zeta(x), \zeta(y))$ belongs to $\partial^2\mathscr{F}$ whenever $x, y \in \partial\Gamma$ are distinct.

Labourie[7] has shown that an important class of representations admits equivariant mappings into Furstenberg's boundary. Consider Σ a closed orientable surface of genus $g \geq 2$ and say that a representation $\pi_1(\Sigma) \to \mathrm{PSL}(d,\mathbb{R})$ is *Fuchsian* if it factors as

$$\pi_1(\Sigma) \to \mathrm{PSL}(2,\mathbb{R}) \to \mathrm{PSL}(d,\mathbb{R})$$

where $\mathrm{PSL}(2,\mathbb{R}) \to \mathrm{PSL}(d,\mathbb{R})$ is the unique irreducible linear action $\mathrm{PSL}(2,\mathbb{R}) \curvearrowright \mathbb{R}^d$ (modulo conjugation by $\mathrm{PSL}(d,\mathbb{R})$) and $\pi_1(\Sigma) \to \mathrm{PSL}(2,\mathbb{R})$ is co-compact. A *Hitchin component* of $\mathrm{PSL}(d,\mathbb{R})$ is a connected component of

$$\hom(\pi_1(\Sigma), \mathrm{PSL}(d,\mathbb{R})) = \{\text{morphisms } \rho : \pi_1(\Sigma) \to \mathrm{PSL}(d,\mathbb{R})\}$$

containing a Fuchsian representation.

Theorem (Labourie[7]). *A representation in a Hitchin component of* $\mathrm{PSL}(d,\mathbb{R})$ *is hyperconvex.*

Denote $C(X)$ the set of real continuous functions on some space X, and $C^*(X)$ its dual. The main result of this work, inspired in Roblin[15], is the following theorem :

Theorem A (See section §5)**.** *Let $\rho : \Gamma \to G$ be a Zariski dense hyperconvex representation, then there exist positive numbers, h and c, and a probability μ on $\partial\Gamma$, such that*

$$ce^{-ht} \sum_{\gamma : d_X(o, \rho(\gamma)\cdot o) \leq t} \delta_{\gamma_-} \otimes \delta_{\gamma_+} \to \mu \otimes \mu$$

on $C^*(\partial\Gamma \times \partial\Gamma)$ as t goes to infinity.

As first shown by Margulis[10] in negative curvature, in order to obtain a counting theorem a mixing property of a well chosen dynamical system is sufficient. In compact manifolds of negative curvature the geodesic flow plays this role. In infinite co-volume, for example for convex co-compact groups, one should restrict the geodesic flow to its non-wandering set, also known as the convex heart of the group.

When Δ is a lattice in higher rank Eskin-McMullen[5] use the mixing property of the *Weyl chamber flow* to prove the counting result previously mentioned:

Let K be a maximal compact subgroup of G τ the Cartan involution on $\mathfrak{g}$ for which the set $\operatorname{fix}\tau$ is K's Lie algebra, consider $\mathfrak{p} = \{v \in \mathfrak{g} : \tau v = -v\}$ and $\mathfrak{a}$ a maximal abelian subspace contained in $\mathfrak{p}$.

Denote $\mathfrak{a}^+ \subset \mathfrak{a}$ a Weyl chamber and M the centralizer of $e^{\mathfrak{a}}$ on K, then the Weyl chamber flow is the action

$$\Delta \backslash G/M \curvearrowleft e^{\mathfrak{a}^+}.$$

The mixing property of this action when Δ is a lattice is shown by Howe-Moore[6].

In this article we show a mixing property of the Weyl chamber flow for hyperconvex representations. Before stating the result let us recall the Patterson-Sullivan theory on higher rank.

Denote $a : G \to \mathfrak{a}^+$ for the Cartan projection (following the Cartan decomposition $G = Ke^{\mathfrak{a}^+}K$) and chose a norm $\| \ \|$ on $\mathfrak{a}$ invariant under the Weyl group. If $\| \ \|$ is induced by a G-invariant Riemannian metric on X then for every $g \in G$ one has $\|a(g)\| = d_X(o, go)$.

Hence one is interested in understanding the growth of

$$\#\{g \in \Delta : \|a(g)\| \leq t\}$$

when $t \to \infty$ for a given subgroup Δ of G. Given an open cone $\mathscr{C} \subset \mathfrak{a}$ consider the exponential growth rate

$$h_{\mathscr{C}} := \limsup_{s \to \infty} \frac{\log \#\{g \in \Delta : a(g) \in \mathscr{C} \text{ with } \|a(g)\| \leq s\}}{s}$$

The *growth indicator function*, introduced by Quint[11], is then the function $\psi_\Delta : \mathfrak{a} \to \mathbb{R} \cup \{-\infty\}$ defined as

$$\psi_\Delta(v) := \|v\| \inf\{h_{\mathscr{C}} : \ \mathscr{C} \text{ open cone with } v \in \mathscr{C}\}.$$

Remark that ψ_Δ is homogeneous and does not depend on the norm chosen.

Quint[11] then shows the following theorem, recall that Benoist[1] has introduced the *limit cone* $\mathscr{L}_\Delta$, of a subgroup Δ, as the closed cone on $\mathfrak{a}^+$ generated by $\{\lambda(g) : g \in \Delta\}$, where $\lambda : G \to \mathfrak{a}^+$ is the Jordan projection.

Theorem 1.1 (Quint[11])**.** *Let Δ be a Zariski dense discrete subgroup of G Then ψ_Δ is concave and upper semi-continuous, the set*

$$\{v \in \mathfrak{a} : \psi_\Delta(v) > -\infty\}$$

is the limit cone $\mathscr{L}_\Delta$ of Δ. ψ_Δ is non negative on $\mathscr{L}_\Delta$ and positive on its interior.

The growth indicator function plays the role in higher rank of the critical exponent in negative curvature. Denote $\sigma : G \times \mathscr{F} \to \mathfrak{a}$ the *Busemann cocycle* introduced by Quint[12]: The set $\mathscr{F}$ is K-homogeneous with stabilizer M, then $\sigma : G \times \mathscr{F} \to \mathfrak{a}$ is defined to verify the following equation

$$gk = l \exp(\sigma(g, kM))n$$

following Iwasawa's decomposition of $G = Ke^{\mathfrak{a}}N$, where N is the unipotent radical of P.

Theorem 1.2 (Quint[12])**.** *Let Δ be a Zariski dense discrete subgroup of G. Then for each functional φ tangent to ψ_Δ in a direction interior to the limit cone, there exists a probability ν_φ on $\mathscr{F}$, supported on the limit set L_Δ, such that for every $g \in \Delta$ one has*

$$\frac{dg_*\nu_\varphi}{d\nu_\varphi}(x) = e^{-\varphi\sigma(g^{-1},x)}.$$

Remark that if $\mathrm{i} : \mathfrak{a}^+ \to \mathfrak{a}^+$ is the *opposition involution* then $\psi_\Delta \mathrm{i} = \psi_\Delta$, since $a(\mathrm{i}\, g) = a(g^{-1})$. Hence, if $\varphi \in \mathfrak{a}^*$ is tangent to ψ_Δ, so is $\varphi \mathrm{i}$. Thus, in higher rank Patterson-Sullivan's measures come in pairs.

As in negative curvature, one can use these measures to construct invariant measures for the Weyl chamber flow. Consider the action of G on $\partial^2\mathscr{F} \times \mathfrak{a}$ via Busemann's cocycle

$$g(x, y, v) = (gx, gy, v - \sigma(g, y)).$$

The stabilizer of a point is isomorphic to M and we get thus an identification $G/M = \partial^2\mathscr{F} \times \mathfrak{a}$. This is called *Hopf's parametrization* of G.

Using Tits[20]'s representations of G one can define a Gromov product

$$\mathscr{G}_\Pi : \partial^2\mathscr{F} \to \mathfrak{a}$$

(see page 20) in order to trivialize Busemann's cocycle on $\partial^2\mathscr{F}$ i.e. for every $g \in G$ and $(x, y) \in \partial^2\mathscr{F}$

$$\mathscr{G}_\Pi(gx, gy) - \mathscr{G}_\Pi(x, y) = -(\mathrm{i}\,\sigma(g, x) + \sigma(g, y)).$$

For a given $\varphi \in \mathfrak{a}^*$ tangent to ψ_Δ, the measure

$$e^{-\varphi(\mathscr{G}_\Pi(\cdot,\cdot))}\nu_{\varphi\circ\mathrm{i}} \otimes \nu_\varphi \otimes \mathrm{Leb}_{\mathfrak{a}}$$

is thus Δ invariant and $\mathfrak{a}$ invariant. Denote χ_φ the measure induced on the quotient $\Delta\backslash G/M$, we call this measure *the Bowen-Margulis measure* for φ by analogy with the negative curvature setting. Its support is the set

$$\Delta\backslash\partial^2\,\mathrm{L}_\Delta \times \mathfrak{a}$$

where $\partial^2\,\mathrm{L}_\Delta = \mathrm{L}_\Delta^2 \cap \partial^2\mathscr{F}$. An important contrast thought is that χ_φ should have infinite total mass.

We then show the following mixing property for hyperconvex representations inspired in the work of Thirion[19] that shows an analogue mixing property for ping-pong groups:

Theorem B (Theorem 4.18)**.** *Let $\rho : \Gamma \to G$ be a Zariski dense hyperconvex representation, $\varphi \in \mathfrak{a}^*$ tangent to ψ_Δ in the direction u_φ. Assume also that u_φ belongs to the interior of the limit cone. Then there exists some positive constant κ such that for any two continuous functions $f_0, f_1 : \rho(\Gamma)\backslash G/M \to \mathbb{R}$ with compact support one has*

$$(2\pi t)^{(\mathrm{rank}(G)-1)/2}\chi_\varphi(f_0 \cdot f_1 \circ e^{tu_\varphi}) \to \kappa\chi_\varphi(f_0)\chi_\varphi(f_1)$$

when $t \to \infty$.

In section §2 we recall results on Hölder cocycles from A.S.[17], of particular interest is the reparametrizing theorem 2.9. This theorem is crucial in understanding the nature of

$$\rho(\Gamma)\backslash\partial^2\,\mathrm{L}_{\rho(\Gamma)} \times \mathfrak{a}$$

when $\rho : \Gamma \to G$ is hyperconvex (proposition 3.2). In section §3 we study vector cocycles and prove a general mixing property that will imply theorem B. This is shown in section §4. In the last section we prove theorem A.

Acknowledgments

I would like to thank Jean-François Quint for proposing me to study the orbital counting problem for Hitchin representations. I'm also thankful for fruitful discussions with him.

2 Real Hölder cocycles

Reparametrizations

Let X be a compact metric space and $\phi_t : X \circlearrowleft$ a continuous flow on X without fixed points.

Definition 2.1. A *translation cocycle* over ϕ_t is a function $\kappa : X \times \mathbb{R} \to \mathbb{R}^d$ that verifies the following two conditions:

- For every $x \in X$ and $t, s \in \mathbb{R}$ one has
$$\kappa(x, t+s) = \kappa(\phi_s(x), t) + \kappa(x, s),$$
- For every $t \in \mathbb{R}$ $\kappa(\cdot, t)$ is Hölder continuous with exponent independent of t and with bounded constant when t is bounded.

The standard example of a translation cocycle is by considering some Hölder continuous function $f : X \to \mathbb{R}^d$ and setting $\kappa_f : X \times \mathbb{R} \to \mathbb{R}^d$ as

$$\kappa_f(x, t) = \int_0^t f\phi_s(x)ds, \tag{1}$$

if t is positive, and $\kappa_f(x, t) := -\kappa_f(\phi_t x, -t)$ for t negative.

Two translation cocycles κ_1 and κ_2 are said to be *Livšic cohomologous* if there exists a continuous $V : X \to \mathbb{R}$ such that for all $x \in X$ and $t \in \mathbb{R}$ one has

$$\kappa_1(x, t) - \kappa_2(x, t) = V(\phi_t x) - V(x).$$

We say that some function $U : X \to \mathbb{R}^d$ is C^1 *in the flow's direction* if for every $x \in X$ the function $t \mapsto U(\phi_t(x))$ is of class C^1 and the function

$$x \mapsto \left.\frac{\partial}{\partial t}\right|_{t=0} U(\phi_t(x))$$

is continuous. Hence, if two translation cocycles κ_f and κ_g are Livšic cohomologous then the function V is C^1 in the flow's direction and

$$f(x) - g(x) = \left.\frac{\partial}{\partial t}\right|_{t=0} V(\phi_t(p)).$$

For a periodic orbit τ of ϕ_t we denote the $p(\tau)$ for its period. If κ is a translation cocycle then the *period* of κ for τ is defined as

$$L_\kappa(\tau) = \kappa(x, p(\tau))$$

for any $x \in \tau$. It is clear that $L_\kappa(\tau)$ does not depend on the point $x \in \tau$ chosen, and that the set of periods is a cohomological invariant of κ.

Assume now that $f : X \to \mathbb{R}$ is positive, then since X is compact f has a positive minimum and $\kappa_f(x, \cdot)$ is an increasing homeomorphism of $\mathbb{R}$. We then have an inverse $\alpha_f : X \times \mathbb{R} \to \mathbb{R}$ that verifies

$$\alpha_f(x, \kappa_f(x, t)) = \kappa_f(x, \alpha_f(x, t)) = t \tag{2}$$

for every $(x, t) \in X \times \mathbb{R}$.

Definition 2.2. The *reparametrization* of ϕ_t by f is the flow $\psi_t : X \circlearrowleft$ defined as $\psi_t(x) := \phi_{\alpha_f(x,t)}(x)$. If f is Hölder continuous we shall say that ψ_t is a Hölder reparametrization of ϕ_t.

Remark 2.1. When two continuous functions $f, g : X \to \mathbb{R}_+^*$ are Livšic cohomologous the reparametrization of ϕ_t by f is conjugated to the reparametrization by g, i.e. there exists a homeomorphism $h : X \to X$ such that for all $p \in X$ and $t \in \mathbb{R}$

$$h(\psi_t^f p) = \psi_t^g(hp).$$

If m is a ϕ_t-invariant probability on X and ψ_t is the reparametrization of ϕ_t by f, then the probability m' defined by $dm'/dm(\cdot) = f(\cdot)/m(f)$ is ψ_t-invariant. In particular, if τ is a periodic orbit of ϕ_t then it is also periodic for ψ_t and the new period is

$$\int_\tau f = \int_0^{p(\tau)} f(\phi_s(x))ds$$

for any $x \in \tau$.

Denote $\mathcal{M}^{\phi_t}$ the set of ϕ_t-invariant probabilities. The *pressure* of a continuous function $f : X \to \mathbb{R}$ is defined as

$$P(\phi_t, f) = \sup_{m \in \mathcal{M}^{\phi_t}} h(\phi_t, m) + \int_X f dm.$$

A probability m such that the supremum is attained is called an *equilibrium state* of f. An equilibrium state for the potential $f \equiv 0$ is called a probability if maximal entropy and its entropy is called the topological entropy of ϕ_t, denoted $h_{\mathrm{top}}(\phi_t)$.

Lemma 2.1 (§2 of A.S.[17])**.** *Consider* $\psi_t : X \circlearrowleft$ *the reparametrization of* $\phi_t : X \circlearrowleft$ *by* $f : X \to \mathbb{R}_+^*$. *Assume* $h := h_{top}(\psi_t) < \infty$, *then the bijection* $m \mapsto m'$ *induces a bijection between equilibrium states of* $-hf$ *and probabilities of maximal entropy of* ψ_t.

Anosov flows and Markov codings

Assume from now on that X is a compact manifold and that the flow $\phi_t : X \circlearrowleft$ is C^1 . We say that ϕ_t is *Anosov* if the tangent bundle of X splits as a sum of three $d\phi_t$-invariant bundles

$$TX = E^s \oplus E^0 \oplus E^u,$$

and there exist positive constants C and c such that: E^0 is the direction of the flow and for every $t \geq 0$ one has: for every $v \in E^s$

$$\|d\phi_t v\| \leq Ce^{-ct}\|v\|,$$

and for every $v \in E^u$ $\|d\phi_{-t} v\| \leq Ce^{-ct}\|v\|$.

We need the following classical result of Livšic[9]:

Theorem 2.2 (Livšic[9]). *Let $\phi_t : X \circlearrowleft$ be an Anosov flow and $\kappa : X \times \mathbb{R} \to \mathbb{R}^d$ a translation cocycle. If $L_\kappa(\tau) = 0$ for every periodic orbit τ, then there exists $V : X \to G$ Hölder continuous such that*

$$\kappa(x,t) = V(\phi_t x) - V(x).$$

As shown by the next lemma, one can always chose a translation cocycle of the form κ_f in the same cohomology class of a given translation cocycle κ.

Lemma 2.3. *Let $\kappa : X \times \mathbb{R} \to \mathbb{R}^d$ be a translation cocycle then there exists $f : X \to \mathbb{R}^d$ Hölder continuous such that the cocycles κ et κ_f are cohomologous.*

Proof. Fix some $C > 0$ and consider the cocycle

$$\kappa^C(x,t) = \frac{1}{C}\int_0^C \kappa(\phi_s(x), t)ds.$$

κ^C and κ are cohomolgous since they have the same periods. One easily checks that $\kappa^C(\cdot, t)$ is of class C^1 in the flow's direction and thus κ^C is the integral of some function on the orbits of ϕ_t. ☐

The following lemma is extremely useful.

Lemma 2.4 (§3 of A.S.[17]). *Consider $f : X \to \mathbb{R}$ Hölder continuous such that*

$$\frac{1}{p(\tau)}\int_\tau f > k$$

for some positive k and every periodic orbit τ of ϕ_t. Then f is Livšic cohomologous to a positive Hölder function.

In order to study ergodic theory of Anosov flows, Bowen[3] and Ratner[14] have introduced the notion of Markov coding.

Definition 2.3. We shall say that the triplet (Σ, π, r) is a *Markov coding* for $\phi_t : X \circlearrowleft$ if Σ is an irreducible subshit of finite type, $\pi : \Sigma \to X$ and $r : \Sigma \to \mathbb{R}_+^*$ are Hölder continuous and the function $\pi_r : \Sigma \times \mathbb{R} \to X$ defined as

$$\pi_r(x,t) = \phi_t \pi(x)$$

verifies the following conditions:

i) π_r is surjective and Hölder,

ii) let $\sigma : \Sigma \circlearrowleft$ be the shift and let $\hat{r} : \Sigma \times \mathbb{R} \circlearrowleft$ be defined as $\hat{r}(x,t) = (\sigma x, t - r(x))$, then π_r is $\hat{r}$-invariant,

iii) $\pi_r : \Sigma \times \mathbb{R}/\hat{r} \to X$ is bounded-to-one and injective on a residual set which is of full measure for every ergodic invariant measure of total support (for σ_t^f),

iv) consider the translation flow $\sigma_t^r : \Sigma \times \mathbb{R}/\hat{r} \circlearrowleft$ then $\pi_r \sigma_t^r = \phi_t \pi_r$.

Remark 2.2. If $\phi_t : X \circlearrowleft$ admits a Markov coding then it has a unique probability of maximal entropy and the function $\pi_r : \Sigma \times \mathbb{R}/\hat{r} \to X$ is an isomorphism between the probabilities of maximal entropy of σ_t^r and that of ϕ_t. In particular the topological entropy of ϕ_t coincides with that of σ_t^r.

We will use the following corollary of A.S.[17]:

Corollary 2.5 (§2 of A.S.[17]). *Let Γ be a co-compact group of isometries of a complete simply connected manifold of negative curvature $\widetilde{M}$. Let $\phi_t : \Gamma\backslash T^1\widetilde{M} \circlearrowleft$ be the geodesic flow and $\psi_t : \Gamma\backslash T^1\widetilde{M} \circlearrowleft$ be a Hölder reparametrization of ϕ_t. Then ψ_t admits a Markov coding (Σ, π, r). The group generated by the periods of the suspension flow σ_t^r is dense in $\mathbb{R}$.*

Hölder cocycles on $\partial\Gamma$

Let Γ be a torsion free co-compact group of isometries of a complete simply connected negatively curved manifold $\widetilde{M}$. The group Γ is hyperbolic and the visual boundary of $\widetilde{M}$ is identified with the boundary of the group. We will now focus on Hölder cocycles on $\partial\Gamma$.

Definition 2.4. A *Hölder cocycle* is a function $c : \Gamma \times \partial\Gamma \to \mathbb{R}$ such that

$$c(\gamma_0\gamma_1, x) = c(\gamma_0, \gamma_1 x) + c(\gamma_1, x)$$

for any $\gamma_0, \gamma_1 \in \Gamma$ and $x \in \partial\Gamma$, and where $c(\gamma, \cdot)$ is a Hölder map for every $\gamma \in \Gamma$ (the same exponent is assumed for every $\gamma \in \Gamma$).

Recall that the dynamics of each $\gamma \in \Gamma$ on $\partial\Gamma$ is of type north-south: γ has two fixed points, γ_+ (an attractor) and γ_-, and the basin of attraction of γ_+ is $\partial\Gamma - \{\gamma_-\}$.

Given a Hölder cocycle c we define the *periods* of c as the numbers

$$\ell_c(\gamma) := c(\gamma, \gamma_+)$$

where γ_+ is the attractive fixed point of γ in $\Gamma - \{e\}$. The cocycle property implies that the period of an element γ only depends on its conjugacy class $[\gamma] \in [\Gamma]$.

Two cocycles c and c' are said to be cohomologous if there exists a Hölder function $U : \partial\Gamma \to \mathbb{R}$ such that for all $\gamma \in \Gamma$ one has

$$c(\gamma, x) - c'(\gamma, x) = U(\gamma x) - U(x).$$

One easily deduces from the definition that the set of periods of a cocycle is a cohomological invariant. We shall use the following theorem of Ledrappier[8] which relates Hölder cocycles with Hölder potentials on $T^1\widetilde{M}$.

Theorem 2.6 (Ledrappier[8], page 105)**.** *For each Hölder cocycle c there exists a Hölder continuous Γ-invariant function $F_c : T^1\widetilde{M} \to \mathbb{R}$ such that for every $\gamma \in \Gamma$ one has*

$$\ell_c(\gamma) := c(\gamma, \gamma_+) = \int_\gamma F_c.$$

A change in the cohomology class of c produces a change in the Livšic cohomology class of F_c.

We are interested in cocycles with positive periods, i.e. $\ell_c(\gamma) > 0$ for every $\gamma \in \Gamma$. The *exponential growth rate* of such cocycle is

$$h_c := \limsup_{s\to\infty} \frac{\log \#\{[\gamma] \in [\Gamma] : \ell_c(\gamma) \leq s\}}{s} \in (0, \infty]$$

where $[\gamma]$ is the conjugacy class of γ. (It is consequence of Ledrappier's work that a Hölder cocycle c with positive periods verifies $h_c > 0$).

Lemma 2.7 (§2 of A.S.[17])**.** *Let $c : \Gamma \times \partial\Gamma \to \mathbb{R}$ be a Hölder cocycle with positive periods and finite exponential growth rate, then the function F_c is Livšic cohomologous to a positive function.*

If c has finite exponential growth rate then, following Patterson's construction, Ledrappier[8] shows the existence of a *Patterson-Sullivan* probability μ over $\partial\Gamma$ of cocycle $h_c c$, that is to say, μ verifies

$$\frac{d\gamma_*\mu}{d\mu}(x) = e^{-h_c c(\gamma^{-1},x)}.$$

Theorem 2.8 (Ledrappier[8] page 102)**.** *Let c be a Hölder cocycle with positive periods. Then c has finite exponential growth rate h_c if and only if there exists a Patterson-Sullivan probability of cocycle $h_c c$. If this is the case, the Patterson-Sullivan probability is unique.*

Let $\overline{c}$ be a cocycle such that $\ell_{\overline{c}}(\gamma) = \ell_c(\gamma^{-1})$, $\overline{c}$ is called *a dual cocycle of c*. Set $\partial^2\Gamma$ to be the set of pairs $(x, y) \in \partial\Gamma \times \partial\Gamma$ such that $x \neq y$. We shall say that a function $[\cdot,\cdot] : \partial^2\Gamma \to \mathbb{R}$ is a *Gromov product* for a pair of dual cocycles $\{c, \overline{c}\}$ if for every $\gamma \in \Gamma$ and $(x, y) \in \partial^2\Gamma$ one has

$$[\gamma x, \gamma y] - [x, y] = -(\overline{c}(\gamma, x) + c(\gamma, y)).$$

Remark 2.3. The existence of these objects for a given Hölder cocycle is consequence of Ledrappier's theorem 2.6. See §2 of A.S.[17] for details

Denote by μ and $\overline{\mu}$ the Patterson-Sullivan probabilities associated to c and $\overline{c}$ respectively. The following theorem is crucial in undertanding the Weyl chamber flow:

Theorem 2.9 (The reparametrizing theorem, A.S.[17])**.** *Let c be a Hölder cocycle with positive periods such that h_c is finite. Then:*

1. *the action of* Γ *in* $\partial^2\Gamma \times \mathbb{R}$ *via* c,

   $$\gamma(x, y, s) = (\gamma x, \gamma y, s - c(\gamma, y)),$$

   is proper and co-compact. Moreover, the translation flow $\psi_t : \Gamma\backslash\partial^2\Gamma \times \mathbb{R} \circlearrowleft$

   $$\psi_t(x, y, s) = (x, y, s - t)$$

   is conjugated to a Hölder reparametrization of the geodesic flow on $\Gamma\backslash T^1\widetilde{M}$*. The conjugating map is also Hölder continuous. The topological entropy of* ψ_t *is* h_c*.*

2. *The measure*

   $$e^{-h_c[\cdot,\cdot]}\overline{\mu} \otimes \mu \otimes ds$$

   on $\partial^2\Gamma \times \mathbb{R}$ *induces on the quotient* $\Gamma\backslash\partial^2\Gamma \times \mathbb{R}$ *the measure of maximal entropy of* ψ_t*.*

3 The action $V \curvearrowright \Gamma\backslash\partial^2\Gamma \times V$

Consider a finite dimensional vector space V and fix some Hölder cocycle $c : \Gamma \times \partial\Gamma \to V$ with values on V. The periods of c are the vectors

$$\ell_c(\gamma) := c(\gamma, \gamma_+) \text{ for } \gamma \in \Gamma.$$

We call $\mathscr{L}_c$ the smallest closed cone of V that contains the vectors $\ell_c(\gamma)$ for every $\gamma \in \Gamma$. The *dual cone* of $\mathscr{L}_c$ is the set of linear functionals positive on this cone:

$$\mathscr{L}_c^* = \{\varphi \in V^* : \varphi|\mathscr{L}_c \geq 0\}.$$

For $\varphi \in \mathscr{L}_c^*$ denote c_φ for the real cocycle $\varphi \circ c$ and h_φ its exponential growth rate:

$$h_\varphi := \limsup_s \frac{\log \#\{[\gamma] : \varphi(\ell_c(\gamma)) \leq s\}}{s}.$$

A direct consequence of the reparametrizing theorem 2.9 is the following.

Corollary 3.1. *If there exists* $\varphi \in \mathscr{L}_c^*$ *such that* h_φ *is finite then the action* $\Gamma \curvearrowright \partial^2\Gamma \times V$ *via* c,

$$\gamma(x, y, v) = (\gamma x, \gamma y, v - c(\gamma, y)),$$

is properly discontinuous.

Proof. One applies theorem 2.9 to the cocycle c_φ. □

We want to understand the topological nature of $\Gamma\backslash\partial^2\Gamma \times V$ together with the action of V. We also need to keep track of a natural measure on $\Gamma\backslash\partial^2\Gamma \times V$ related to c.

Assume from now on the existence of $\varphi \in \mathscr{L}_c^*$ with h_φ finite. Fix also a dual cocycle $\overline{c}$ of c and a vector Gromov product $[\cdot,\cdot] : \partial^2\Gamma \to V$ for the pair $\{c, \overline{c}\}$.

Denote μ_φ and $\overline{\mu}_\varphi$ for the Patterson-Sullivan probabilities of cocycle $h_\varphi c_\varphi$ and $h_\varphi \overline{c}_\varphi$ respectively. The function

$$[\cdot,\cdot]_\varphi := \varphi \circ [\cdot,\cdot]$$

is a Gromov product for the pair $\{c_\varphi, \overline{c}_\varphi\}$. The measure induced on the quotient by

$$e^{-h_\varphi[\cdot,\cdot]_\varphi}\overline{\mu}_\varphi \otimes \mu_\varphi \otimes \mathrm{Leb}_V$$

where Leb_V is a Lebesgue measure on V, is denoted Ω_φ and called the *Bowen-Margulis measure* of the pair $\{c, \overline{c}\}$ for the functional φ.

Definition 3.1. Fix some Γ-invariant Hölder continuous function $F : T^1\widetilde{M} \to V$ given by Ledrappier's theorem 2.6 for the cocycle c. For $\varphi \in \mathscr{L}_c^*$ denote m_φ for the equilibrium state on T^1M of the function $-h_\varphi\varphi \circ F$. Then, the *dual direction* of $h_\varphi\varphi$ is the direction in V given by the vector

$$\int F dm_\varphi.$$

The dual direction of φ is the same for any functional in $\mathbb{R}_+\varphi$ and is independent on the choice of F since a change in F's cohomology (Livšic class) does not change the value of the integral $\int F dm$ for any ϕ_t-invariant probability measure.

Remark that since $h_\varphi < \infty$ lemma 2.7 implies that the function $\varphi(F)$ is cohomologous to a positive function and thus we can choose some u_φ in the dual direction of φ such that $\varphi(u_\varphi) = 1$. Consider the flow $\omega_{tu_\varphi} : \Gamma\backslash\partial^2\Gamma \times V \circlearrowleft$ defined as the quotient of

$$(x, y, v) \mapsto (x, y, v - tu_\varphi).$$

Proposition 3.2. *Assume there exists $\varphi \in \mathscr{L}_c^*$ such that h_φ is finite. Then there exists a Hölder reparametrization of the geodesic flow $\psi_t : T^1M \circlearrowleft$ and a Hölder function $f : T^1M \to \ker\varphi$ with*

$$\int_{T^1M} f dm' = 0.$$

for ψ_t's probability of maximal entropy m', and a Hölder continuous function

$$\overline{E} : \Gamma\backslash\partial^2\Gamma \times V \to T^1M \times \ker\varphi$$

that conjugates flow ω_{tu_φ} with the flow $\widetilde{\psi}_t : T^1M \times \ker\varphi \circlearrowleft$ defined as

$$\widetilde{\psi}_t(p, v_0) = (\psi_t(p), v_0 - \int_0^t f\psi_s(p)ds).$$

$\overline{E}$ also conjugates the action of $\ker\varphi$ and is an isomorphism (modulo a constant) between the measures Ω_φ and $m' \otimes \mathrm{Leb}_{\ker\varphi}$.

Proof. Consider the function $\widetilde{\varphi} : \partial^2\Gamma \times V \to \partial^2\Gamma \times \mathbb{R}$ given by

$$\widetilde{\varphi} : (x, y, v) \mapsto (x, y, \varphi(v)).$$

The action of Γ on $\partial^2\Gamma \times V$ is read, via $\widetilde{\varphi}$, as

$$\gamma(x, y, t) = (\gamma x, \gamma y, t - c_\varphi(\gamma, y))$$

and the measure Ω_φ descends on $\partial^2\Gamma \times \mathbb{R}$ as

$$e^{-h_\varphi[\cdot,\cdot]_\varphi}\overline{\mu}_\varphi \otimes \mu_\varphi \otimes \mathrm{Leb}_\mathbb{R} .$$

Since h_φ is finite the reparametrizing theorem 2.9 applies and thus the action of Γ on $\partial^2\Gamma \times \mathbb{R}$ via c_φ is properly discontinuous. Moreover there exists $E : \Gamma\backslash\partial^2\Gamma \times \mathbb{R} \to \Gamma\backslash T^1M$ Hölder continuous which conjugates the translation flow with ψ_t and the image of the measure (induced in the quotient by)

$$e^{-h_\varphi[\cdot,\cdot]_\varphi}\overline{\mu}_\varphi \otimes \mu_\varphi \otimes \mathrm{Leb}_\mathbb{R}$$

is ψ_t's measure of maximal entropy. We can assume (via a cohomology choice) that $\varphi(F)$ is strictly positive (lemma 2.4) and thus the flow ψ_t is the reparametrization of the geodesic ϕ_t by $\varphi(F)$. We then have that ψ_t's probability of maximal entropy is m'_φ (recall that m_φ is the equilibrium state of $-h_\varphi\varphi(F)$, and use lemma 2.1).

Considering the composition $E \circ \widetilde{\varphi} : \Gamma\backslash\partial^2 \times V \to \Gamma\backslash T^1M$ one finds that, for every $u \in V$,

$$(x, y, v - u) \mapsto \psi_{\varphi(u)}(E(x, y, \varphi(v))).$$

In particular the flow ω_{tu_φ} descends on T^1M as ψ_t.

The action of the abelian group $\ker\varphi$ on $\partial^2\Gamma \times V$ commutes with Γ's action and preserves the fibers $\widetilde{\varphi}^{-1}(x, y, t)$ of $\widetilde{\varphi}$. This action induces thus an action of $\ker\varphi$ on the quotient and one finds that

$$E \circ \widetilde{\varphi} : \Gamma\backslash\partial^2\Gamma \times V \to T^1M$$

is a vector bundle with fiber $\ker\varphi$, and that the group $\ker\varphi$ acts on $\Gamma\backslash\partial^2\Gamma \times V$ preserving the fibers, this is

$$\ker\varphi \to \Gamma\backslash\partial^2\Gamma \times V \to T^1M$$

is a principal bundle with group $\ker\varphi$. Using the zero section of a vector bundle, and the action of $\ker\varphi$, one can trivialize this principal bundle. Hence, $\Gamma\backslash\partial^2\Gamma \times V$ is (Hölder) isomorphic to $T^1M \times \ker\varphi$.

Set $\Psi_t : T^1M \times \ker\varphi \circlearrowleft$ to be the corresponding flow to $\omega_{tu_\varphi} : \Gamma\backslash\partial^2\Gamma \times V \circlearrowleft$ via this last identification. Since w_{tu_φ} commutes with the action of $\ker\varphi$, the same occurs for Ψ_t and thus one can write

$$\Psi_t(p, v_0) = (\psi_t(p), v_0 - \kappa(p, t))$$

where $\kappa : T^1M \times \mathbb{R} \to \ker\varphi$ is a translation cocycle over ψ_t. Lemma 2.3 implies the existence of a Hölder continuous function $f : T^1M \to \ker\varphi$ such that the cocycles κ and $(x,t) \mapsto \int_0^t f\psi_s(x)ds$ are cohomologous (for the dynamics ψ_t). The flow Ψ_t is then conjugated to the flow

$$\widetilde{\psi_t}(x,v) = (\psi_t(x), v - \int_0^t f\psi_s(x)ds).$$

We remark that the image of the measure Ω_φ on $T^1M \times \ker\varphi$ is a measure that descends to T^1M as m'_φ and is $\ker\varphi$-invariant. This measure is then $m'_\varphi \otimes \mathrm{Leb}_{\ker\varphi}$.

It remains to check that $\int_{T^1M} f dm'_\varphi = 0$. In order to do this recall that $\varphi(u_\varphi) = 1$ and that u_φ is co-linear to the vector $\int F dm_\varphi$, moreover

$$\int F dm_\varphi = u_\varphi \int \varphi(F) dm_\varphi.$$



Figure 1: If $p \in T^1M$ belongs to the periodic orbit associated to $[\gamma]$, the translation on the fiber $\ker\varphi$, by the flow $\widetilde{\psi}_t$ at the returning time, is given by $\ell_c^0(\gamma)$.

Consider $F = F_0 + \varphi(F)u_\varphi$ following the decomposition $V = \ker\varphi \oplus \mathbb{R}u_\varphi$. One obtains thus that

$$\int_{T^1M} F_0 dm_\varphi = 0.$$

For $\gamma \in \Gamma$ let $\ell_c^0(\gamma)$ be the projection of the period $\ell_c(\gamma)$ on $\ker\varphi$ following the decomposition $V = \ker\varphi \oplus \mathbb{R}u_\varphi$. One has that

$$\ell_c^0(\gamma) = \int_{[\gamma]} F_0\phi_s ds.$$

The fact that $\widetilde{\psi}_t$ is conjugated to the flow w_{tu_φ} implies that

$$\ell_c^0(\gamma) = \int_{[\gamma]'} f\psi_s ds = \int_{[\gamma]} f\phi_s\varphi(F\phi_s)ds$$

(see figure) where $[\gamma]'$ is the periodic of ψ_t associated to $[\gamma]$. Livšic's theorem 2.2 then implies that the functions F_0 and $f\varphi(F)$ are cohomologous (for the dynamics ϕ_t), and thus

$$0 = \int F_0 dm_\varphi = \int f\varphi(F)dm_\varphi = \int f dm'_\varphi.$$

This finishes the proof.

□

Mixing properties of V's action

Now that we have a good description of $\Gamma\backslash\partial^2\Gamma \times V$ together with V's action we can use Markov codings, and a theorem of Thirion[19] for suspensions of sub-shifts of finite type, to prove the following mixing property:

For $v_0 \in \ker\varphi$ and $t \in \mathbb{R}^+$ denote $w_t^{v_0} : \Gamma\backslash\partial^2\Gamma \times V \circlearrowleft$ defined as the induced in the quotient of

$$(x, y, v) \mapsto (x, y, v - tu_\varphi - \sqrt{t}v_0).$$

Theorem 3.3. *Let $c : \Gamma \times \partial\Gamma \to V$ be a Hölder cocycle such that the group generated by its periods is dense in V. Suppose there exists a linear functional $\varphi \in \mathscr{L}_c^*$ such that $h_\varphi < \infty$. Then $w_t^{v_0} : \Gamma\backslash\partial^2\Gamma \times V \circlearrowleft$ verifies: given two compactly supported continuous functions $f_0, f_1 : \Gamma\backslash\partial^2\Gamma \times V \to \mathbb{R}$ then*

$$(2\pi t)^{(\dim V - 1)/2} \int_{\Gamma\backslash\partial^2\Gamma\times V} f_0(z)f_1(w_t^{v_0}z)d\Omega_\varphi(z)$$

converges, when $t \to \infty$, to

$$k(v_0)\Omega_\varphi(f_0)\Omega_\varphi(f_1)$$

for some constant $k(v_0)$.

Proof. Applying proposition 3.2 we get a reparametrization of the geodesic flow $\psi_t : T^1M \circlearrowleft$ together with a Hölder continuous function $f : T^1M \to \ker\varphi$ and $\widetilde{E} : \Gamma\backslash\partial^2\Gamma \times V \to T^1M \times \ker\varphi$ that conjugates:

- the action of $\ker\varphi$ on $\Gamma\backslash\partial^2\Gamma \times V$ and on $T^1M \times \ker\varphi$,

- the flow ω_{tu_φ} with

$$(x,v) \mapsto (\psi_t(x), v - \int_0^t f\psi_s(x)ds)$$

Hence we will study mixing properties of

$$t \cdot (x,v) \mapsto (\psi_t(x), v - \int_0^t f\psi_s ds - \sqrt{t}v_0).$$

□

Consider (Σ_A, π, r) a Markov coding for $\psi_t : T^1M \circlearrowleft$ (corollary 2.5). There exists an equilibrium state of the shift $\sigma : \Sigma_A \circlearrowleft$, denoted ν_φ, corresponding to the measure m'_φ via the Markov coding, i.e. for every measurable function $G : T^1M \to \mathbb{R}$ one has

$$\int_{T^1M} G dm'_\varphi = \int_{\Sigma_A} \int_0^{r(x)} G(\psi_s \pi(x)) ds d\nu_\varphi.$$

Define $K : \Sigma_A \to V$ given by

$$K(x) = \int_0^{rx} f\psi_s(\pi x)ds + r(x)u_\varphi$$

and $\hat{K} : \Sigma_A \times V \circlearrowleft$ as $\hat{K}(x,v) = (\sigma x, v - K(x))$.

Lemma 3.4. *The function* $\underline{\pi} : \Sigma_A \times V \to T^1M \times \ker\varphi$, *defined as*

$$\begin{aligned}\underline{\pi}(x,v) &= (\psi_{\varphi(v)}(\pi x), v - \varphi(v)u_\varphi - \int_0^t f\psi_s(\pi x)ds) \\ &= \widetilde{\psi}_{\varphi(v)}(\pi x, v - \varphi(v)u_\varphi)\end{aligned}$$

is $\hat{K}$*-invariant and induces measurable isomorphism between* $\nu_\varphi \otimes \mathrm{Leb}_V$ *and* $m'_\varphi \otimes \mathrm{Leb}_{\ker\varphi}$.

Proof. The proof follows directly from the properties of Markov codings. □

Hence, the flow $\widetilde{\psi}_t$ is measurably conjugated to the translation flow on $\Sigma_A \times V/\hat{K}$ in the direction given by the vector

$$u_\varphi = \int_{\Sigma_A} K d\nu_\varphi.$$

Moreover, this conjugation also conjugates the action of $\ker\varphi$ on $T^1M \times \ker\varphi$ and on $\Sigma_A \times V$. The theorem 3.3 is thus consequence of the following theorem due to Thirion[19].

Theorem 3.5 (Thirion[19])**.** *Let* Σ_A *be a sub-shift of finite type* $K:\Sigma_A\to V$ *Hölder continuous function and* ν *an equilibrium state. Assume there exists* $\varphi\in V^*$ *such that* $\varphi(K)$ *is cohomologous to a positive function. We also suppose that the group generated by the periods of* K *is dense in* V*. Set*

$$\tau:=\int_{\Sigma_A}Kd\nu\in V.$$

Then the translation flow $\Sigma_A\times V/\hat{K}$ *in the direction of* τ *verifies: given two compactly supported continuous functions* $f_0,f_1:\Sigma_A\times V/\hat{K}\to\mathbb{R}$ *and* $v_0\in\ker\varphi$ *one has*

$$(2\pi t)^{(\dim V-1)/2}\int_{\Sigma_A\times V/\hat{K}}f_0(x,v)f_1(x,v-t\tau-\sqrt{t}v_0)d\nu\otimes\mathrm{Leb}_V$$

converges when $t\to\infty$*, to*

$$k(v_0)\int_{\Sigma_A\times V/\hat{K}}f_0d\nu\otimes\mathrm{Leb}_V\int_{\Sigma_A\times V/\hat{K}}f_1d\nu\otimes\mathrm{Leb}_V$$

where $k(v_0)$ *is some constant depending on* v_0*.*

4 Convex representations and the Weyl chamber flow

We are now interested in studying representations $\Gamma\to G$ admitting equivariant maps to $\mathscr{F}_\theta=G/P_\theta$ where P_θ is some parabolic group of G. Recall that Γ is the fundamental group of some closed negatively curved manifold M.

Consider a real semi-simple algebraic group G. Let K be a maximal compact subgroup of G τ the Cartan involution on $\mathfrak{g}$ for which the set $\operatorname{fix}\tau$ is K's Lie algebra, consider $\mathfrak{p}=\{v\in\mathfrak{g}:\tau v=-v\}$ and $\mathfrak{a}$ a maximal abelian subspace contained in $\mathfrak{p}$.

Let Σ be the roots of $\mathfrak{a}$ on $\mathfrak{g}$, Σ^+ a system of positive roots on Σ and Π the set of simple roots associated to the choice Σ^+. To each subset θ of Π one associates a parabolic subgroup P_θ of G whose Lie algebra is, by definition,

$$\mathfrak{p}_\theta=\mathfrak{a}\oplus\bigoplus_{\alpha\in\Sigma^+}\mathfrak{g}_\alpha\oplus\bigoplus_{\alpha\in\langle\Pi-\theta\rangle}\mathfrak{g}_{-\alpha}$$

where $\langle\theta\rangle$ is the set of positive roots generated by θ and

$$\mathfrak{g}_\alpha=\{w\in\mathfrak{g}:[v,w]=\alpha(v)w\ \forall v\in\mathfrak{a}\}.$$

Every parabolic subgroup of G is conjugated to a unique P_θ.

Set W to be the Weyl group of Σ and note $w_0:\mathfrak{a}\to\mathfrak{a}$ the biggest element in W, w_0 is the unique element in W that sends $\mathfrak{a}^+$ to $-\mathfrak{a}^+$. The *opposition involution* $\mathrm{i}:\mathfrak{a}\to\mathfrak{a}$ is the defined as $\mathrm{i}:=-w_0$.

Fix from now on $\theta \subset \Pi$ a subset of simple roots of G and write $\mathscr{F}_\theta$ for G/P_θ. We consider also $P_{\mathrm{i}\,\theta}$, the parabolic group associated to

$$\mathrm{i}\,\theta := \{\alpha \circ \mathrm{i} : \alpha \in \theta\}.$$

The set $\mathscr{F}_{\mathrm{i}\,\theta} \times \mathscr{F}_\theta$ possesses a unique open G-orbit, which we will denote $\partial^2\mathscr{F}_\theta$.

Definition 4.1. We say that a representation $\rho : \Gamma \to G$ is θ-*convex* if it admits two Hölder continuous equivariant maps $\xi : \partial\Gamma \to \mathscr{F}_\theta$ and $\eta : \partial\Gamma \to \mathscr{F}_{\mathrm{i}\,\theta}$ such that whenever $x \neq y$ in $\partial\Gamma$ one has that the pair $(\xi(x), \eta(y))$ belongs to $\partial^2\mathscr{F}_\theta$.

A Π-convex representation (i.e., when the set θ is the full set of simple roots and thus the parabolic group P_Π is minimal) is called *hyperconvex.* The set $\mathscr{F}_\Pi$ is the Furstenberg boundary $\mathscr{F}$ of G's symmetric space.

An element $g \in G$ is said to be *proximal on* $\mathscr{F}_\theta$ if it has an attracting fixed point on $\mathscr{F}_\theta$. This points is noted g_+^θ. One finds that such g also has a fixed point g_-^θ on $\mathscr{F}_{\mathrm{i}\,\theta}$ wich is the attractor for g^{-1} on $\mathscr{F}_{\mathrm{i}\,\theta}$ and every point $x \in \mathscr{F}_\theta$ such that the pair (g_-^θ, x) belongs to $\partial^2\mathscr{F}_\theta$ verifies $g^n x \to g_+^\theta$. The point g_-^θ is called the repelling hyperplane of g.

Lemma 4.1 (§3 of A.S.[16])**.** *Let $\rho : \Gamma \to G$ be a Zariski dense θ-convex representation, then for every $\gamma \in \Gamma$, $\rho(\gamma)$ is proximal on $\mathscr{F}_\theta$, $\xi(\gamma_+)$ is its attracting fixed point and $\eta(\gamma_-)$ is the repelling hyperplane.*

The equivariant functions ξ and η of the definition are then unique since attracting points γ_+ are dense in $\partial\Gamma$. The image of ξ (reps. η) is then the limit set of $\rho(\Gamma)$ on $\mathscr{F}_\theta$ (resp. $\mathscr{F}_{\mathrm{i}\,\theta}$).

An element $g \in G$ is said to be purely loxodromic if its conjugated to an element on $Me^{\mathfrak{a}^{++}}$ where $\mathfrak{a}^{++}$ is the interior of the Weyl chamber.

The corollary implies, in particular, that hyperconvex representations are purely loxodromic. Nevertheless one has the following stronger result, recall that by definition, the limit cone of a subgroup is closed.

Lemma 4.2 (§3 of A.S.[16])**.** *The limit cone of a Zariski dense hyperconvex representation is contained in the interior of the Weyl chamber $\mathfrak{a}^+$.*

Busemann cocycle

Given a θ-convex representation $\rho : \Gamma \to G$ there is a natural Hölder (vector) cocycle on the boundary of Γ that appears, for which we need *Busemann's cocycle* on G introduced by Quint[12]:

The set $\mathscr{F}$ is K-homogeneous with stabilizer M, where K is a maximal compact subgroup of G, one then defines $\sigma_\Pi : G \times \mathscr{F} \to \mathfrak{a}$ to verify the following equation

$$gk = l \exp(\sigma_\Pi(g, kM))n$$

following Iwasawa's decomposition of $G = Ke^{\mathfrak{a}}N$, where N is the unipotent radical of P.

In order to obtain a cocycle only depending on the set $\mathscr{F}_\theta$ (and G) one considers

$$\mathfrak{a}_\theta := \bigcap_{\alpha\in\Pi-\theta} \ker\alpha$$

the lie algebra of the center of the reductive group $P_\theta \cap \overline{P_\theta}$, where $\overline{P_\theta}$ is the opposite parabolic group of P_θ. Consider also $p_\theta : \mathfrak{a} \to \mathfrak{a}_\theta$ the only projection invariant under the group $W_\theta = \{w \in W : w(\mathfrak{a}_\theta) = \mathfrak{a}_\theta\}$.

Remark 4.1. One easily verifies the following relation: $p_{\mathrm{i}\,\theta} = \mathrm{i}\, p_\theta\, \mathrm{i}$.

Quint[12] then shows the following lemma:

Lemma 4.3 (Lemma 6.1 of Quint[12])**.** *The function $p_\theta \circ \sigma_\Pi$ factors trough a function $\sigma_\theta : G \times \mathscr{F}_\theta \to \mathfrak{a}_\theta$. σ_θ verifies the cocycle relation: for every $g, h \in G$ and $x \in \mathscr{F}_\theta$ one has*

$$\sigma_\theta(gh, x) = \sigma_\theta(g, hx) + \sigma_\theta(h, x).$$

The cocycle one naturally associates to a θ-convex representation is then $\beta_\theta : \Gamma \times \partial\Gamma \to \mathfrak{a}_\theta$ defined as

$$\beta_\theta(\gamma, x) = \sigma_\theta(\rho(\gamma), \xi(x)).$$

Every $g \in G$ decomposes into $g = g_s g_u$ where g_s is semisimple and g_u is unipotent. The *Jordan projection* $\lambda : G \to \mathfrak{a}^+$ is defined as $g_s = me^{\lambda(g)}$ for some $m \in M$.

Lemma 4.4. *Let $\rho : \Gamma \to G$ be Zariski dense and θ-convex. Then for every $\gamma \in \Gamma$ the period $\beta_\theta(\gamma, \gamma_+)$ of β_θ is*

$$\beta_\theta(\gamma, \gamma_+) = p_\theta(\lambda(\rho\gamma)) := \lambda_\theta(\rho\gamma).$$

Proof. The proof follows exactly as lemma 7.5 of A.S.[17]. □

Consider the Hölder cocycle

$$\overline{\beta_\theta} := \mathrm{i}\,\beta_{\mathrm{i}\,\theta}.$$

Since $\mathrm{i}\, p_\theta\, \mathrm{i} = p_{\mathrm{i}\,\theta}$ and $\mathrm{i}(\lambda(g)) = \lambda(g^{-1})$ for every $g \in G$ one obtains the following corollary:

Corollary 4.5. *The pair $\{\beta_\theta, \overline{\beta_\theta}\}$ is a pair of dual cocycles.*

Consider the cone $\mathscr{L}_\rho^\theta$, this is the closed cone generated by

$$\{\lambda_\theta(\rho\gamma) : \gamma \in \Gamma\}$$

and consider its dual cone

$$\mathscr{L}_\rho^{\theta^*} := \{\varphi \in \mathfrak{a}_\theta^* : \varphi|\mathscr{L}_\rho^\theta \geq 0\}.$$

Since $\mathscr{L}_\rho^\theta$ is contained in $p_\theta(\mathfrak{a}^+)$ it does not contain any line, the dual cone is thus $\mathscr{L}_\rho^{\theta^*}$ with non empty interior.

Lemma 4.6. *Let $\rho : \Gamma \to G$ be a Zariski dense θ-convex representation and consider some φ in the interior of the dual cone $\mathscr{L}_\rho^{\theta^*}$ then the cocycle $\varphi \circ \beta_\theta : \Gamma \times \partial\Gamma \to \mathbb{R}$ has finite exponential growth rate.*

Proof. The proof of the lemma follows exactly as lemma 7.7 of A.S.[17]. ☐

Applying corollary 3.1 to the cocycle β_θ one directly obtains:

Corollary 4.7. *Let $\rho : \Gamma \to G$ be Zariski dense and θ-convex, then the action $\Gamma \curvearrowright \partial^2\Gamma \times \mathfrak{a}_\theta$ is properly discontinuous.*

Even thought we will not use it on this paper we remark that lemma 4.6 together with corollary 4.1 of A.S.[17] imply the following counting result:

Corollary 4.8. *Let $\rho : \Gamma \to G$ be Zariski dense and θ-convex, and consider φ in the interior of $\mathscr{L}_\rho^{\theta^*}$, then there exists h_φ such that*

$$h_\varphi t e^{-h_\varphi t} \#\{[\gamma] \in [\Gamma] : \varphi(\lambda_\theta(\rho\gamma)) \leq t\} \to 1$$

when $t \to \infty$.

Gromov product

The purpose of this section is to define a Gromov product for the pair $\{\beta_\theta, \overline{\beta_\theta}\}$. We begin with the following result of Tits[20].

Proposition 4.9 (Tits[20]). *For each $\alpha \in \Pi$ there exists a finite dimensional proximal irreducible representation $\Lambda_\alpha : G \to \mathrm{PGL}(V_\alpha)$ such that the highest weight χ_α of Λ_α is an integer multiple of the fundamental weight ω_α. Moreover, any other weight of Λ_α is of the form*

$$\chi_\alpha - \alpha - \sum_{\beta \in \Pi} n_\beta \beta$$

with $n_\beta \in \mathbb{N}$.

Fix some θ and consider some $\alpha \in \theta$. Consider also $\Lambda_\alpha : G \to \mathrm{PGL}(V_\alpha)$ a representation given by Tits's proposition. Since Λ_α is proximal, one obtains an equivariant mapping $\xi_\alpha : \mathscr{F}_\theta \to \mathbb{P}(V_\alpha)$.

The highest weight of the dual representation $\Lambda_\alpha^* : G \to \mathbb{P}(V_\alpha^*)$ is $\chi_\alpha \,\mathrm{i}$, one thus obtains an equivariant mapping $\eta_\alpha : \mathscr{F}_{\mathrm{i}\,\theta} \to \mathbb{P}(V_\alpha^*)$. Moreover, the pair $(x, y) \in \partial^2\mathscr{F}_\theta$ verifies

$$\eta_\alpha(x)|\xi_a(y) \neq 0.$$

We define the *Gromov product* $\mathscr{G}_\theta : \partial^2\mathscr{F}_\theta \to \mathfrak{a}_\theta$ as follows: Since $\{\omega_\alpha|\mathfrak{a}_\theta : a \in \theta\}$ is a basis of $\mathfrak{a}_\theta^*$ the same occurs for $\{\chi_\alpha|\mathfrak{a}_\theta\}_{\alpha\in\theta}$. The element $\mathscr{G}_\theta(x, y)$ is thus determined by $\chi_\alpha(\mathscr{G}_\theta(x, y))$ for every $\alpha \in \theta$.

Consider a scalar product on V_α such that $\Lambda_\alpha(K)$ is orthogonal and $\exp \mathfrak{a}$ is symmetric. The euclidean norm $\| \ \|_\alpha$ induced by this scalar product verifies

$$\log \|\Lambda_\alpha(g)\|_\alpha = \chi_\alpha(a(g))$$

where $a : G \to \mathfrak{a}^+$ is the Cartan projection. We then define $\chi_\alpha(\mathscr{G}_\theta(x,y))$ as

$$\chi_\alpha(\mathscr{G}_\theta(x,y)) := \log \frac{|\varphi(v)|}{\|\varphi\|_\alpha \|v\|_\alpha}$$

for some $\varphi \in \eta_\alpha(x)$ and $v \in \xi_\alpha(y)$.

Lemma 4.10. *For every $g \in G$ and $x, y \in \partial^2 \mathscr{F}_\theta$ one has*

$$\mathscr{G}_\theta(gx, gy) - \mathscr{G}_\theta(x,y) = -(\mathrm{i}\,\sigma_{\mathrm{i}\,\theta}(g,x) + \sigma_\theta(g,y)).$$

Proof. The lemma follows from the formula

$$\log \frac{|\varphi \circ g^{-1}(gv)|}{\|\varphi \circ g^{-1}\| \|gv\|} - \log \frac{|\varphi(v)|}{\|\varphi\|\|v\|} = -\log \frac{\|g\varphi\|}{\|\varphi\|} + \log \frac{\|gv\|}{\|v\|},$$

for a norm on a vector space V, every $g \in \mathrm{PGL}(V)$ and $(\varphi, v) \in \mathbb{P}(V^*) \times \mathbb{P}(V) - \{(\varphi, v) \in \mathbb{P}(V^*) \times \mathbb{P}(V) : \varphi(v) = 0\}$, together with the definition of $\mathscr{G}_\theta$ and the fact that for every $\alpha \in \theta$

$$\chi_\alpha(\sigma(g,x)) = \log \frac{\|\Lambda_\alpha(g)v\|_\alpha}{\|v\|_\alpha}$$

for $v \in \xi_\alpha(x)$ (lemma 6.4 of Quint[12]). □

The following corollary is immediate.

Corollary 4.11. *Let $\rho : \Gamma \to G$ be a Zariski dense θ-convex representation and consider φ in the interior of the dual cone ${\mathscr{L}_\rho^\theta}^*$, then the function $[\cdot,\cdot]_\varphi : \partial^2\Gamma \to \mathbb{R}$ defined as*

$$[x,y]_\varphi = \varphi \circ \mathscr{G}_\theta(\eta(x), \xi(y))$$

is a Gromov product for the pair of dual cocycles $\{\varphi(\beta_\theta), \varphi(\overline{\beta_\theta})\}$.

We need the following theorem of Benoist[2]:

Theorem 4.12 (Benoist[2])**.** *Consider Δ a Zariski dense subgroup of G, then the group generated by $\{\lambda(g) : g \in \Delta\}$ is dense in $\mathfrak{a}$.*

Recall that the Bowen-Margulis measure for φ is the measure Ω_φ on $\Gamma\backslash\partial^2\Gamma \times \mathfrak{a}_\theta$ defined as the induced in the quotient of

$$e^{-h_\varphi[\cdot,\cdot]_\varphi}\overline{\mu_\varphi} \otimes \mu_\varphi \otimes \mathrm{Leb}_{\mathfrak{a}_\theta}$$

where μ_φ and $\overline{\mu_\varphi}$ are the Patterson-Sullivan probabilities of cocycles $\varphi(\beta_\theta)$ and $\varphi(\overline{\beta_\varphi})$ respectively. Benoist[2]'s theorem guarantees the missing hypothesis of theorem 3.3 and we obtain thus the following.

Theorem 4.13. *Let $\rho : \Gamma \to G$ be a Zariski dense θ-convex representation, and consider φ in the interior of the cone ${\mathscr{L}_\rho^\theta}^*$. Denote u_φ^θ the dual direction of φ. Then there exists some positive constant κ such that for any two continuous functions $f_0, f_1 : \Gamma\backslash\partial^2\Gamma \times \mathfrak{a}_\theta \to \mathbb{R}$ with compact support one has*

$$(2\pi t)^{(\dim \mathfrak{a}_\theta - 1)/2}\Omega_\varphi(f_0 \cdot f_1 \circ \omega_{tu_\varphi^\theta}) \to \kappa\Omega_\varphi(f_0)\Omega_\varphi(f_1)$$

when $t \to \infty$.

Remark 4.2. When θ is not the full set of simple roots, the last theorem does not give a mixing property of the Weyl chamber flow of $\rho(\Gamma)$, nevertheless it may turn out to be useful since

$$\rho(\Gamma)\backslash\partial^2 \mathrm{L}_{\rho(\Gamma)} \times\mathfrak{a}$$

fibers over

$$\Gamma\backslash\partial^2\Gamma \times \mathfrak{a}_\theta$$

(where $\mathrm{L}_{\rho(\Gamma)}$ is the limit set of $\rho(\Gamma)$ on $\mathscr{F}$).

The growth indicator function

Denote $a : G \to \mathfrak{a}^+$ for the Cartan projection (following the Cartan decomposition $G = Ke^{\mathfrak{a}^+}K$), $X = G/K$ for G's symmetric space and fix a norm $\| \ \|$ on $\mathfrak{a}$ invariant under the Weyl group. If $\| \ \|$ is induced by a G-invariant Riemannian metric on X then one has $\|a(g)\| = d_X(o, go)$ where $o = [K]$ for every $g \in G$.

Hence one is interested in understanding the growth

$$h_\Delta^{\| \ \|} := \limsup_{s\to\infty} \frac{\log \#\{g \in \Delta : \|a(g)\| \leq s\}}{s}$$

for a given subgroup Δ of G. Given an open cone $\mathscr{C} \subset \mathfrak{a}$ we consider the exponential growth rate

$$h_{\mathscr{C}} := \limsup_{s\to\infty} \frac{\log \#\{g \in \Delta : a(g) \in \mathscr{C} \text{ with } \|a(g)\| \leq s\}}{s},$$

the *growth indicator function*, introduced by Quint[11], is then the function $\psi_\Delta : \mathfrak{a} \to \mathbb{R} \cup \{-\infty\}$ defined as

$$\psi_\Delta(v) := \|v\| \inf\{h_{\mathscr{C}} : \ \mathscr{C} \text{ open cone with } v \in \mathscr{C}\}.$$

Remark that ψ_Δ does not depend on the norm chosen. Quint[11] then shows the following theorem.

Theorem 4.14 (Quint[11]). *Let Δ be a Zariski dense discrete subgroup of G Then ψ_Δ is concave and upper semi-continuous, the set*

$$\{v \in \mathfrak{a} : \psi_\Delta(v) > -\infty\}$$

is the limit cone $\mathscr{L}_\Delta$ of Δ. ψ_Δ is non negative on $\mathscr{L}_\Delta$ and positive on its interior.

One then retrieves the exponential growth rate for a given norm as

$$\sup_{v\in\mathfrak{a}-\{0\}} \frac{\psi_\Delta(v)}{\|v\|} = h_\Delta^{\| \ \|}.$$

Assume from now on that $\| \ \|$ comes from a G-invariant Riemannian metric on X and set $h_\Delta = h_\Delta^{\| \ \|}$. Since ψ_Δ is concave, and the balls of $\| \ \|$ are strictly convex, one obtains a unique direction u_Δ on the limit cone $\mathscr{L}_\Delta$ such that the supremum

$$\sup \frac{\psi_\Delta}{\|\ \|}$$

is attained. This is called *the growth direction* of Δ. One remarks that the number of elements in $a(\Delta)$ that lie outside a given open cone containing u_Δ has exponential growth rate strictly smaller than h_Δ.

Denote ∂X for X's visual boundary and $\overline{X} = X \cup \partial X$. Consider the function $\pi : \mathscr{F} \times \mathfrak{a}^+ \to \partial X$ defined as

$$\pi(kM, v) = \pi_v(kM) = \lim_{t\to\infty} ke^{tv} \cdot o$$

(recall that $\mathscr{F} = K/M$ and $o = [K]$). The following lemma is direct:

Lemma 4.15. *Let Δ be a Zariski dense subgroup of G, u_Δ its growth direction, and consider a continuous function $f : \overline{X} \to \mathbb{R}$ whose support does not intersect the $\pi_{u_\Delta}(\mathscr{F})$, then*

$$e^{-h_\Delta t} \sum_{g\in\Delta : d_X(o,go)\leq t} f(g\cdot o) \to 0$$

when t goes to infinity.

Proof. The lemma follows directly from the definition of the growth direction u_Δ. ☐

If one considers a linear functional $\varphi \in \mathfrak{a}^*$ such that $\varphi(v) \geq \psi_\Delta(v)$, then $\|\varphi\| \geq h_\Delta^{\|\ \|}$. One is thus interested in the set

$$D_\Delta = \{\varphi \in \mathfrak{a}^* : \varphi \geq \psi_\Delta\}.$$

Since ψ_ρ is homogeneous the set D_Δ is convex, hence it has a point closest to the origin, Θ_Δ. This linear form is *the growth form* of Δ. We remark that Θ_Δ is tangent to the growth indicator ψ_Δ in the growth direction u_Δ.

Fix from now on $\rho : \Gamma \to G$ a Zariski dense hyperconvex representation. The cone $\mathscr{L}_\rho^\Pi$ is the limit cone $\mathscr{L}_\rho$ of $\rho(\Gamma)$.

In A.S.[16] we prove the following result:

Theorem 4.16 (A.S.[16])**.** *The growth indicator ψ_ρ of a Zariski dense hyperconvex representation ρ is strictly concave, analytic on the interior of the limit cone and with vertical tangent on its boundary. If φ belongs to the interior of the cone $\mathscr{L}_\rho^*$ then $h_\varphi\varphi$ is tangent to ψ_ρ in the dual direction of φ.*

Remark 4.3. Hence for a Zariski dense hyperconvex representation, the growth direction u_ρ is the dual direction of the growth form.

The set $\mathscr{F}_\Pi = \mathscr{F}$ is Furstenberg's boundary of G's symmetric space. Denote $\zeta : \partial\Gamma \to \mathscr{F}$ for ρ's equivariant map. The image $\zeta(\partial\Gamma)$ is the limit set $\mathrm{L}_{\rho(\Gamma)}$ and thus

$$\zeta \times \zeta : \partial^2\Gamma \to \partial^2\,\mathrm{L}_{\rho(\Gamma)}$$

is a ρ-equivariant Hölder homeomorphism.

Corollary 4.17. *Let $\rho : \Gamma \to G$ be a Zariski dense hyperconvex representation. For each φ tangent to ψ_ρ there exists a unique Patterson-Sullivan measure for φ denoted ν_φ. Moreover, ζ induces an isomorphism between μ_φ and ν_φ.*

Consequently the function

$$\zeta \times \zeta \times \mathrm{id} : \partial^2\Gamma \times \mathfrak{a} \to \partial^2 \mathrm{L}_{\rho(\Gamma)} \times \mathfrak{a}$$

induces a ρ-equivariant isomorphism between the Bowen-Margulis measures of φ on each side:

$$\Omega_\varphi \leftrightarrow \chi_\varphi.$$

Hence, theorem 4.13 together with remark 4.3 imply the following mixing property of the Weyl chamber flow. Recall that the rank of G is the dimension of $\mathfrak{a}$.

Theorem 4.18. *Let $\rho : \Gamma \to G$ be a Zariski dense hyperconvex representation, consider Θ its growth form and u_ρ is growth direction. Fix some $v_0 \in \ker\Theta$, then there exists some positive constant $\kappa(v_0)$ such that for any two continuous functions $f_0, f_1 : \rho(\Gamma)\backslash G/M \to \mathbb{R}$ with compact support one has*

$$(2\pi t)^{(\mathrm{rank}(G)-1)/2}\chi_\Theta(f_0 \cdot f_1 \circ e^{tu_\rho + \sqrt{t}v_0}) \to \kappa(v_0)\chi_\Theta(f_0)\chi_\Theta(f_1)$$

when $t \to \infty$.

5 The orbital counting problem

In this section we show theorem A, for which we need a result of Thirion[18].

Theorem 5.1 (Thirion[18])**.** *Let Δ be a Zariski dense discrete subgroup of G and Θ its growth form. Assume its growth direction u_Θ belongs to the interior of the Weyl chamber $\mathfrak{a}^+$ and that the following bound holds:*

$$\{g \in \Delta : d_X(o, go) \leq t\} \leq Ke^{\|\Theta\|t}. \tag{3}$$

If the Weyl chamber flow satisfies: given two compactly supported continuous functions $f_0, f_1 : \Delta\backslash G/M \to \mathbb{R}$ one has

$$(2\pi t)^{(\mathrm{rank}(G)-1)/2}\chi_\Theta(f_0 \cdot f_1 \circ e^{tu_\Theta + \sqrt{t}v_0}) \to \kappa(v_0)\chi_\Theta(f_0)\chi_\Theta(f_1)$$

when $t \to \infty$, for some constant $\kappa(v_0) > 0$, then there exists $c > 0$ such that

$$ce^{-\|\Theta\|t} \sum_{g\in\Delta:d(o,go)\leq t} \delta_{g^{-1}o} \otimes \delta_{go} \to \pi^*_{u_\Theta}\nu_\Theta \otimes \pi^*_{u_\Theta}\nu_\Theta,$$

when $t \to \infty$ in $C^*(\overline{X}^2)$ where ν_Θ is the Patterson-Sullivan measure of the growth form Θ.

Proof of theorem A. Fix $\rho : \Gamma \to G$ a Zariski dense hyperconvex representations. We want to show that

$$ce^{-ht} \sum_{\gamma : d_X(o, \rho(\gamma) \cdot o) \leq t} \delta_{\gamma_-} \otimes \delta_{\gamma_+} \to \mu_\Theta \otimes \mu_\Theta$$

on $C^*(\partial\Gamma \times \partial\Gamma)$ as t goes to infinity, where μ_Θ is the Patterson-Sullivan probability on $\partial\Gamma$ of cocycle $\Theta \circ \beta_\Pi : \Gamma \times \partial\Gamma \to \mathbb{R}$, Θ being the growth form of $\rho(\Gamma)$. We will use the theorem of Thirion to obtain a convergence on $\overline{X}^2$ and then take the measures to $\partial\Gamma \times \partial\Gamma$.

We must thus verify the hypothesis of Thirion's theorem 5.1:

- the growth direction belongs to the interior of the Weyl chamber: this is direct consequence of lemma 4.2,
- there exists $C > 0$ such that $\#\{\gamma \in \Gamma : d_X(o, \rho(\gamma)o) \leq t\} \leq Ce^{\|\Theta\| t}$, this is shown in corollary 7.15 of A.S.[17],
- the mixing property of the Weyl chamber flow is the statement of theorem 4.18.

We thus obtain convergence

$$ce^{-\|\Theta\| t} \sum_{\gamma \in \Gamma : d_X(o, \rho(\gamma)o) \leq t} \delta_{\rho(\gamma^{-1})o} \otimes \delta_{\rho(\gamma)o} \to \pi^*_{u_\Theta} \nu_\Theta \otimes \pi^*_{u_\Theta} \nu_\Theta,$$

when $t \to \infty$ in $C^*(\overline{X}^2)$ where ν_Θ is the Patterson-Sullivan measure of the growth form Θ of $\rho(\Gamma)$.

Using lemma 4.15 we obtain convergence of

$$ce^{-\|\Theta\| t} \sum_{\gamma \in \Gamma : d_X(o, \rho(\gamma)o) \leq t} \delta_{\pi_{u_\Theta} \zeta(\gamma_-)} \otimes \delta_{\pi_{u_\Theta} \zeta(\gamma_+)} \to \pi^*_{u_\Theta} \nu_\Theta \otimes \pi^*_{u_\Theta} \nu_\Theta,$$

on $C^*((\partial X)^2)$ when $t \to \infty$.

This is to say

$$ce^{-\|\Theta\| t} \sum_{\gamma \in \Gamma : d_X(o, \rho(\gamma)o) \leq t} \pi^*_{u_\Theta} \delta_{\zeta(\gamma_-)} \otimes \pi^*_{u_\Theta} \delta_{\zeta(\gamma_+)} \to \pi^*_{u_\Theta} \nu_\Theta \otimes \pi^*_{u_\Theta} \nu_\Theta,$$

when $t \to \infty$ on $C^*((\partial X)^2)$. Since π_{u_Θ} is an homeomorphism onto its image we obtain the convergence

$$ce^{-\|\Theta\| t} \sum_{\gamma \in \Gamma : d_X(o, \rho(\gamma)o) \leq t} \delta_{\zeta(\gamma_-)} \otimes \delta_{\zeta(\gamma_+)} \to \nu_\Theta \otimes \nu_\Theta,$$

when $t \to \infty$ on $C^*(\mathscr{F}^2)$. The fact that $\zeta_* \nu_\Theta = \mu_\Theta$ (corollary 4.17) finishes the proof. □

References

[1] Y. Benoist. Propriétés asymptotiques des groupes linéaires. *Geom. funct. anal.*, 7(1), 1997.

[2] Y. Benoist. Propriétés asymptotiques des groupes linéaires ii. *Adv. Stud. Pute Math.*, 26, 2000.

[3] R. Bowen. Symbolic dynamics for hyperbolic flows. *Amer. J. Math.*, 95, 1973.

[4] W. Duke, Z. Rudnick, and P. Sarnak. Density of integer points on affine homogeneous varieties.sar. *Duke Math. Jour.*, 71(1), 1993.

[5] A. Eskin and C. McMullen. Mixing, counting and equidistribution in Lie groups. *Duke Math. Jour.*, 71, 1993.

[6] R. Howe and C. Moore. Asymptotic properties of unitary representations. *J. Func. Anal.*, 32, 1979.

[7] F. Labourie. Anosov flows, surface groups and curves in projective space. *Invent. Math.*, 165, 2006.

[8] F. Ledrappier. Structure au bord des variétés à courbure négative. *Séminaire de théorie spectrale et géométrie de Grenoble*, 71, 1994-1995.

[9] A.N. Livšic. Cohomology of dynamical systems. *Math. USSR Izvestija*, 6, 1972.

[10] G. Margulis. Applications of ergodic theory to the investigation of manifolds with negative curvature. *Functional Anal. Appl.*, 3, 1969.

[11] J.-F. Quint. Divergence exponentielle des sous-groupes discrets en rang supérieur. *Comment. Math. Helv.*, 77, 2002.

[12] J.-F. Quint. Mesures de Patterson-Sullivan en rang supérieur. *Geom. funct. anal.*, 12, 2002.

[13] J.-F. Quint. Groupes de Schottky et comptage. *Ann. Inst. Fourier*, 55, 2005.

[14] M. Ratner. Markov partitions for anosov flows on n-dimensional manifolds. *Israel Journal of Math.*, 15(1), 1973.

[15] T. Roblin. *Ergodicité et équidistribution en courbure negative*, volume 95 of *Mémoires de la SMF*. Société mathématique de France, 2003.

[16] A. Sambarino. Hyperconvex representations and exponential growth. Submited, 2011.

[17] A. Sambarino. Quantitative properties of convex representations. Submited. arXiv:1104.4705v1, 2011.

[18] X. Thirion. *Sous-groupes discrets de $SL(d, \mathbb{R})$ et equidistribution dans les espaces symetriques.* PhD thesis, Université de Tours, Juin 2007.

[19] X. Thirion. *Propriétés de mélange du flot des chambres de Weyl des groupes de Ping-Pong*, volume 137 of *Bulletin de la SMF.* Société mathématique de France, 2009.

[20] J. Tits. Représentations linéaires irréductibles d'un groupe réductif sur un corps quelconqe. *J. Reine Angew. Math.*, 247, 1971.

Andrés Sambarino
Laboratoire de Mathématiques
Université Paris Sud,
F-91405 Orsay France,
`andres.sambarino@math.u-psud.fr`